

Numerical Methods for Quasicrystals

Kai Jiang^{1, 2, a)} and Pingwen Zhang^{1, b)}

¹⁾*LMAM, CAPT and School of Mathematical Sciences, Peking University, Beijing 100871, P.R. China*

²⁾*Hunan Key Laboratory for Computation and Simulation in Science and Engineering, Xiangtan University, Xiangtan 411105, P.R. China*

(Dated: 17 December 2012)

A new projection method, implemented in higher-dimensional reciprocal space, is developed to compute quasicrystals. The approach enables us to represent quasicrystals as periodic structures in higher-dimensional space. A proposed projective matrix can project a higher-dimensional periodic structure into a quasicrystal in physical space. Compared with the traditional crystalline approximant method, the projection method overcomes the restrictions of the Simultaneous Diophantine Approximation, and can use periodic boundary conditions conveniently. Meanwhile, the proposed method can efficiently reduce the computational complexity through implementing in a unit cell and using pseudospectral method. By applying the projection method to the Lifshitz-Petrich model, we can compute quasicrystals rather than crystalline approximants, and maintain the rotational symmetry accurately. More significantly, the algorithm can calculate the free energy density to high precision.

^{a)}Electronic mail: kaijiang@math.pku.edu.cn

^{b)}Electronic mail: pzhang@pku.edu.cn

I. INTRODUCTION

As early as the 1890s, the periodic structures (crystals) in three dimensions were determined by 230 space groups based on periodicity, and then the classical crystallography was completed, in which the allowed rotational symmetry is only 1-, 2-, 3-, 4-, 6-fold symmetry. Both the structure determination and the study of physical properties are based on the periodicity which allows the study to be simplified to a unit cell. However, in the 1980s, a forbidden 5-fold symmetry electron diffraction pattern was discovered by Shechtman et al.¹ in a rapid cooled Al-Mn alloy. Later, the term “quasicrystals” appeared for the first time² to describe the non-conventional ordered structures. In an idealized description, quasicrystals have quasiperiodic, rather than periodic, translational order with non-crystallographic symmetry. In the early theory of tilings, the discovery of aperiodic tiling with 5-fold symmetry of the plane by Penrose^{3,4} showed that such well-ordered systems were mathematically possible. Since the original discovery, hundreds of quasicrystals have been reported and confirmed in metallic alloys with 5-, 8-, 10-, 12- fold orientational symmetry^{5,6}. Shechtman also won the Nobel Prize in Chemistry in 2011 for the discovery of quasicrystals. Two decades after the first discovery of quasicrystals in metallic alloys, several soft quasicrystals have been found in soft matter systems. A number of materials successively joined the family of soft quasicrystals, including supramolecular dendrimer or tetraphilic liquid crystals⁷⁻¹⁰, *ABC* star terpolymers¹¹, surfactant-coated metallic nanoparticles¹², and colloidal copolymer micelles in solution¹³. Recently the natural quasicrystals have also been discovered in a khatyrkite-bearing sample¹⁴. To date, quasicrystals are confirmed as a new class of ordered structures.

Two kinds of theoretical approaches have been developed to study quasicrystals, or more generally, aperiodic crystals. The first method is an approximate approach which studies crystalline approximants rather than quasicrystals. The crystalline approximants are periodic structures in which the arrangements of lattices closely approximate the local structures in quasicrystals. Many crystalline approximants related to quasicrystals have also been discovered¹⁵. These approximants may play an important role in describing the local structures of quasicrystals, their formation, stability, and physical properties. The second approach is a direct method to study the quasicrystals or aperiodic crystals in the hyperspace, called the higher-dimensional approach. In this method, a quasiperiodic structure may be viewed as

a periodic structure by extending it into a higher-dimensional space. Its symmetries can be expressed in terms of the conventional point groups and space groups of higher-dimensional periodic crystals^{16,17}. A simple and famous example is the one-dimensional Fibonacci sequence. The sequence is formed by iterative application of the substitution rule : $L \rightarrow LS$, $S \rightarrow L$ to the two-letter alphabet $\{L, S\}$. Then the n -th Fibonacci number F_n satisfies the recurrence relation $F_{n+1} = F_n + F_{n-1}$. The ratio $\lim_{n \rightarrow \infty} F_n/F_{n-1} = \tau = (\sqrt{5} + 1)/2$ is irrational, and is the golden ratio. Therefore the Fibonacci sequence is a quasiperiodic series. From another point of view, as Fig. 1 shows, the one-dimensional Fibonacci sequence




FIG. 1: Fibonacci sequence obtained as the projection of the square lattice inside the window whose width $\sqrt{2}a$ is shown between broken and solid lines with a slope of

$$\alpha = 1/\tau = (\sqrt{5} - 1)/2.$$

can be obtained as a projection from a two-dimensional square-lattice of points. If the slope α is rational, the projected series becomes a periodic chain. In the higher-dimensional description, quasiperiodic structures result from irrational physical-space cuts of appropriate periodic hypercrystal structures. The higher-dimensional approach reveals the hidden structural correlations. To implement the higher-dimensional approach in the direct space one must know the discrete lattice arrangements of higher-dimensional periodic structure. The embedded spaces of d -dimensional quasiperiodic structures are abstract spaces whose dimensions are more than three. The dimensions of the embedded space are dependent on the symmetry of the quasicrystal ($d > 1$)¹⁷. For example, the quasicrystals with 5-, 8-,

10-, and 12-fold symmetry need to be embedded into four-dimensional space. While for the quasiperiodic structures with 7-, 9-, 18-fold symmetry, the dimension of the embedding spaces increases to six. For other symmetries, embedding spaces with even higher dimension will be needed¹⁸. Therefore it can make the method difficult to implement. Another disadvantage of applying the higher-dimensional approach in direct space is that in practical problems one is required to compute the continuous quasiperiodic distribution rather than the discrete quasiperiodic lattice. A convenient way to describe the quasiperiodic structures is in the higher-dimensional reciprocal space¹⁷. By redefining point-group symmetry and space-group symmetry in terms of gauge functions, a broader Fourier-space crystallography¹⁹⁻²¹ was developed to describe quasicrystals.

In this paper, we focus on the development of numerical methods for generating quasicrystals based on the idea of Fourier-space crystallography¹⁹⁻²¹. Quasicrystals are one kind of space-filling structures. The traditional numerical method uses a periodic structure to approximate the quasicrystal²². In other words, those methods compute a crystalline approximant. A natural expectation is that the obtained approximants should approximate quasicrystals as the computational box goes to infinity. The advantage of the approximate method is that using the periodic boundary conditions is convenient. However, the approach cannot obtain quasicrystals exactly unless the computational box is the whole space because of the restriction of the Simultaneous Diophantine Approximation (SDA)²³. We will further explain this point in Sec. II. We then will provide a new numerical method based on the higher-dimensional reciprocal space description^{19,20} to calculate quasicrystals rather than crystalline approximants. The proposed numerical method is able to overcome the restriction of SDA and compute the energy density to high accuracy without periodic approximation.

Although our method is applicable to any model including quasicrystals, for definiteness, we use the Lifshitz-Petrich model²² to demonstrate our algorithm. Before we go further, a short introduction to the Lifshitz-Petrich model is necessary. Lifshitz-Petrich model is a mean-field theory, and its free energy density functional is

$$F[\phi(x, y)] = \frac{1}{V} \int dx dy \left\{ \frac{c}{2} [(\nabla^2 + 1)(\nabla^2 + q^2)\phi]^2 - \frac{\varepsilon}{2} \phi^2 - \frac{\alpha}{3} \phi^3 + \frac{1}{4} \phi^4 \right\}. \quad (1)$$

In Eqn. (1), $\phi(x, y)$ is the order parameter. V is the system volume. q is an irrational number depending on the symmetry. ε is the reduced temperature. $c > 0$ is an energy penalty to

ensure that the principle reciprocal vectors of structures is located on $|\mathbf{k}| = 1$ and $|\mathbf{k}| = q$. $\alpha > 0$ is a phenomenological parameter. For quasicrystals the system volume V should go to infinity since quasicrystals are the space-filling structures without periodicity. For ordered phases including periodic and quasiperiodic ones, the order parameter ϕ^* is the minimum of the free energy density functional, which means

$$\left. \frac{\delta F}{\delta \phi(\mathbf{r})} \right|_{\phi^*} = 0. \quad (2)$$

In order to find the equilibrium state, we introduce a relaxational dynamical to minimize the Lifshitz-Petrich energy functional which yields

$$\frac{\partial \phi}{\partial t} = -\frac{\delta F}{\delta \phi} = -c(\nabla^2 + 1)^2(\nabla^2 + q^2)^2\phi + \varepsilon\phi + \alpha\phi^2 - \phi^3. \quad (3)$$

We choose a semi-implicit scheme to solve the dynamical equation (3):

$$\frac{1}{\Delta t}(\phi_{t+\Delta t} - \phi_t) = \varepsilon\phi_t - c(\nabla^2 + 1)^2(\nabla^2 + q^2)^2\phi_{t+\Delta t} + \alpha(\phi^2)_t - (\phi^3)_t. \quad (4)$$

Δt is the time step size. The specific implementation of the semi-implicit method for different numerical methods will be discussed in Sec. II.

II. NUMERICAL METHODS

A. Crystalline Approximant Method (CAM)

1. Method Description of CAM

Numerical methods designed for periodic structures have been used to study quasicrystals approximately²². Here we call this method the “crystalline approximant method (CAM)”. In order to describe this method, a brief introduction of numerical methods for treating periodic structures is necessary, more details can be found in [24]. For any d -dimensional periodic function $f(\mathbf{r})$, $\mathbf{r} \in \mathcal{R}^d$, the repeated structural unit is called a unit cell. A primitive unit cell, described by d vectors $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_d$, has the smallest possible volume. The Bravais lattice vector is then defined by

$$\mathbf{R} = l_1\mathbf{e}_1 + l_2\mathbf{e}_2 + \dots + l_d\mathbf{e}_d, \quad (5)$$

where $\mathbf{l} = (l_1, l_2, \dots, l_d)$ is a d -dimensional vector with components $l_i \in \mathcal{Z}$. For any $\mathbf{R}$ in the Bravais lattice, the structure is invariant under a lattice translation, i.e., $f(\mathbf{r} + \mathbf{R}) = f(\mathbf{r})$. Given the primitive vectors $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_d)$, the primitive reciprocal vectors $(\mathbf{e}_1^*, \mathbf{e}_2^*, \dots, \mathbf{e}_d^*)$ satisfy the equation

$$\mathbf{e}_i \cdot \mathbf{e}_j^* = 2\pi\delta_{ij}. \quad (6)$$

The reciprocal lattice vector is then specified by

$$\mathbf{k} = k_1\mathbf{e}_1^* + k_2\mathbf{e}_2^* + \dots + k_d\mathbf{e}_d^*, \quad (7)$$

where $k_i \in \mathcal{Z}$. One of the most important properties of the reciprocal lattices is that plane waves $\{e^{i\mathbf{k}\cdot\mathbf{r}}\}$ form a set of basis functions for any function with periodicity of the lattice. The periodic function $f(\mathbf{r})$ on the Bravais lattice can be expanded as

$$f(\mathbf{r}) = \sum_{\mathbf{k}} \hat{f}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}}. \quad (8)$$

For periodic structures, the reciprocal lattice vectors have two important features: $\mathbf{e}_1^*, \mathbf{e}_2^*, \dots, \mathbf{e}_d^*$ are linearly independent; and $k_i \in \mathcal{Z}$, $i = 1, 2, \dots, d$. In the direct space, the lattice vectors have the same properties.

The main idea of CAM is to use periodic structures to approximate quasicrystals. For a d -dimensional quasicrystals, its reciprocal lattice vectors $\mathbf{k}$ can be expressed by d linearly independent reciprocal vectors, $\mathbf{e}_1^*, \mathbf{e}_2^*, \dots, \mathbf{e}_d^*$,

$$\mathbf{k} = p_1\mathbf{e}_1^* + p_2\mathbf{e}_2^* + \dots + p_d\mathbf{e}_d^*. \quad (9)$$

It is important to note that the $p_i \in \mathcal{R}$, that is, $\mathbf{k}$ cannot be represented by linear combinations of $\mathbf{e}_i^*$ with integer-valued coefficients. However, the quasiperiodic function $\phi(\mathbf{r})$ can be expanded as in the following form

$$\phi(\mathbf{r}) = \sum_{\mathbf{k}} \hat{\phi}(\mathbf{k}) e^{i(L\mathbf{k})\cdot\mathbf{r}}, \quad \mathbf{r} \in [0, 2\pi L)^d. \quad (10)$$

If there exists a real number L such that $Lp_i \in \mathcal{Z}$ or Lp_i can be made arbitrarily close to an integer for all $i = 1, 2, \dots, d$ and $\mathbf{k}$, which means

$$|L \cdot (p_1, p_2, \dots, p_d) - ([Lp_1], [Lp_2], \dots, [Lp_d])|_{l^\infty} \rightarrow 0, \quad \text{for all } \mathbf{k}, \quad (11)$$

where $[\cdot]$ rounds the number $\cdot$ to the nearest integer. Then numerical methods designed for periodic structures can be used to treat quasicrystals. Without loss of generality, we can always choose $(1, 0, \dots, 0)$ as one of the primitive reciprocal vectors. Therefore, the real number L becomes an integer. The problem of determining L is a well-known problem called the Simultaneous Diophantine Approximation (SDA) in number theory²³. In view of numerical computability, the integer L must be as small as possible.

From the above description, the condition of SDA must be satisfied for the implementation of CAM. The integer L is dependent on these irrational numbers p_i due to rotational symmetry and the desired precision of the approximation. In the subsequent numerical examples (in Sec. II C), we will find that L increases very quickly as the desired precision becomes higher. This greatly increases the computational cost. For CAM, the computational box should satisfy the condition (11). Therefore, the edge length $D_{\mathbf{k}}$ of computational box is close to $D_{\mathbf{k}} = L \times 2\pi$, or $D = n \cdot L \times 2\pi$, $n \in \mathcal{N}^+$.

2. An Example of 12-fold Rotational Symmetry




FIG. 2: Sets of reciprocal lattice vectors $\{\mathbf{k}\}$ of 12-fold rotational symmetry in 2-dimensional space.

In order to make the above method clearer, we take an example of 12-fold rotational symmetry in two-dimensions to demonstrate this point. A diagram in the reciprocal space is shown in Fig. 2. Let two noncollinear vectors $\mathbf{e}_1^* = (1, 0)$ and $\mathbf{e}_2^* = (0, 1)$ comprise the primitive reciprocal vectors, other reciprocal vectors of the 12-fold case are represented as linear combinations of $\mathbf{e}_1^*$ and $\mathbf{e}_2^*$ with real number coefficients, as shown in Fig. 2. If other

two noncollinear reciprocal vectors are chosen as primitive reciprocal vectors, similar results will emerge as well. The distinct nonzero coefficients for these 12 vectors are 1, $1/2$, and $\sqrt{3}/2$. When applying CAM to this example of 12-fold vectors, we need an integer L such that

$$\left| L \cdot \left(1, \frac{\sqrt{3}}{2}, \frac{1}{2} \right) - \left(L, \left\lceil \frac{\sqrt{3}}{2} L \right\rceil, \left\lceil \frac{1}{2} L \right\rceil \right) \right|_{l^\infty} \rightarrow 0. \quad (12)$$

3. Applying CAM to Lifshitz-Petrich Method

In CAM, we indeed use a crystalline approximant to approximate the quasicrystal. By substituting Eqn (10) into (1), the Lifshitz-Petrich free energy density functional becomes

$$F[\phi] = \frac{1}{2} \sum_{\mathbf{k}_1 + \mathbf{k}_2 = 0} [c(1 - |\mathbf{k}|^2)^2(q^2 - |\mathbf{k}|^2)^2 - \varepsilon] \hat{\phi}(\mathbf{k}_1) \hat{\phi}(\mathbf{k}_2) - \frac{\alpha}{3} \sum_{\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 = 0} \hat{\phi}(\mathbf{k}_1) \hat{\phi}(\mathbf{k}_2) \hat{\phi}(\mathbf{k}_3) + \frac{1}{4} \sum_{\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3 + \mathbf{k}_4 = 0} \hat{\phi}(\mathbf{k}_1) \hat{\phi}(\mathbf{k}_2) \hat{\phi}(\mathbf{k}_3) \hat{\phi}(\mathbf{k}_4). \quad (13)$$

Then we use semi-implicit scheme (4) to minimize the free energy density functional. For CAM, the semi-implicit method becomes

$$\left(\frac{1}{\Delta t} + c(1 - \mathbf{k}^2)^2(q^2 - \mathbf{k}^2)^2 \right) \hat{\phi}_{t+\Delta t}(\mathbf{k}) = \left(\frac{1}{\Delta t} + \varepsilon \right) \hat{\phi}_t(\mathbf{k}) + \alpha(\phi_t^2)(\mathbf{k}) - (\phi_t^3)(\mathbf{k}), \quad (14)$$

where $(\phi_t^m)(\mathbf{k}) = \int d\mathbf{r} \phi^m(\mathbf{r}) e^{-i\mathbf{k}\mathbf{r}}$, $m = 2, 3$. The right terms of the dynamics can be efficiently calculated by the pseudospectral method²⁴. The Laplacian terms are computed in d -dimensional reciprocal space easily, while the convolutions can be calculated efficiently by Fast-Fourier transformation (FFT). Then the computational complexity is $O(N \log N)$ at each time step, N is the number of degrees of freedom.

B. Projection Method (PM)

1. Method Description of PM

As mentioned above, CAM computes the crystalline approximants. However, the approximation method cannot evaluate the free energy density exactly because of the approximate error of SDA. Therefore it is necessary to design a new numerical method that improves the calculation of quasicrystals by avoiding the approximate error of SDA. From Fourier-space crystallography^{17,19,20}, there exists an equivalent n -dimensional representation of a

d -dimensional quasicrystal ($n > d$). In the higher-dimensional description^{17,19,20}, the reciprocal vector $\mathbf{k}$ of a d -dimensional quasicrystal is,

$$\mathbf{k} = h_1 \mathbf{p}_1^* + h_2 \mathbf{p}_2^* + \cdots + h_n \mathbf{p}_n^*, \quad h_i \in \mathcal{Z}, \quad (15)$$

with vectors $\mathbf{p}_i^* \in \mathcal{R}^d$ of rank n . In the following, one should keep in mind that $\mathbf{k} \in \mathcal{R}^d$. We can redefine the selected reciprocal vectors $\mathbf{p}_i^*$ with n components. Then the quasicrystal can be represented by these redefined basic reciprocal vectors with integral coefficients in n -dimensional space. This representation has the advantage, for our purpose, that the d -dimensional quasicrystal is periodic in n -dimensions. In order to make the description more clear, we introduce the n -dimensional reciprocal lattice. Assume that n -dimensional vectors, $\mathbf{b}_1^*, \mathbf{b}_2^*, \dots, \mathbf{b}_n^*$, are the primitive reciprocal vectors of a 1st Brillouin zone in n -dimensional reciprocal space, the reciprocal vector of n -dimensional periodic structure can be expressed as

$$\mathbf{H} = h_1 \mathbf{b}_1^* + h_2 \mathbf{b}_2^* + \cdots + h_n \mathbf{b}_n^*, \quad (16)$$

where the coefficient $h_i \in \mathcal{Z}$, and $\mathbf{H} \in \mathcal{R}^n$. The correspondingly primitive vectors of the direct space $\mathbf{b}_i$ satisfy the dual relationship (6). We want to use the n -dimensional reciprocal vector $\mathbf{H}$ to represent the d -dimensional quasicrystal. Then we can solve a d -dimensional quasicrystal as a periodic structure in n -dimensional space. The key point of implementing the above idea is to provide an operator to project the n -dimensional structure into d -dimensional space.

In order to solve the problem, we propose a novel Fourier expansion for the d -dimensional quasiperiodic function acting d -dimensions

$$g(\mathbf{r}) = \sum_{\mathbf{H}} \hat{g}(\mathbf{H}) e^{i[(\mathcal{S} \cdot \mathbf{H})^T \cdot \mathbf{r}]}. \quad (17)$$

In the expression, $\mathbf{r} \in \mathcal{R}^d$, $\mathbf{H} \in \mathcal{R}^n$, and $\mathcal{S}$ is the *projective matrix* which connects the d -dimensional physical space with the n -dimensional reciprocal space. We note that the two representations (15) and (16) can be used to describe the same quasicrystal. The reciprocal vectors of a d -dimensional quasicrystal can be represented by d -dimensional reciprocal vectors $\mathbf{p}_i^*$ with integral coefficients, and also the integral combinations of extended n -dimensional primitive reciprocal vectors $\mathbf{b}_i^*$. Therefore, we can obtain the projective matrix $\mathcal{S}$ through projecting the n -dimensional reciprocal vectors $\mathbf{b}_i$ into d -dimensional reciprocal

space, i.e., $\mathbf{p}_i = (\mathcal{S} \cdot \mathbf{b})_i$, $i = 1, 2, \dots, n$. The j -th component p_{ij}^* of the projected reciprocal vector $\mathbf{p}_i^*$ can be expressed by $\mathbf{b}_i^*$

$$p_{ij}^* = \sum_{m=1}^n s_{jm} b_{im}^*, \quad j = 1, 2, \dots, d, \quad (18)$$

where b_{im}^* is the m -th component of the i -th n -dimensional reciprocal vector $\mathbf{b}_i^*$. Then these coefficients $s_{jm} \in \mathcal{R}$, $j = 1, 2, \dots, d$, $m = 1, 2, \dots, n$, form the $d \times n$ -order nonzero projective matrix $\mathcal{S}$. For the least computational expense, the dimension n of the extended space must be the smallest determined by the order of the elements of the symmetric group^{17,18}. For example, 5-, 8-, 10-, and 12-fold symmetric quasicrystals, the dimension of embedded space is four. However, 7-, 9-, and 18-fold symmetric quasicrystals must be restricted to six-dimensional space. The projective matrix $\mathcal{S}$ is not unique, which is determined by the selection of reciprocal vectors $\mathbf{b}_i^*$. The represented coefficients of reciprocal vectors are dependent on the selection of primitive reciprocal vectors, however, the reciprocal vectors of a quasicrystal are unique. Therefore, the selection of basic reciprocal vectors $\mathbf{b}_i^*$ as well as the projective matrix $\mathcal{S}$ is irrelevant to the quasicrystal. Considering a periodic structure, the projective matrix $\mathcal{S}$ degenerates to a $d \times d$ -order unit matrix.

Furthermore, PM can be extended to calculate one-dimensional quasicrystals even though the notion of rotational symmetry does not exist in one dimension. If an energy functional has one-dimensional quasiperiodic structures with different incommensurate scales, for example,

$$g(x) = C_0 \sin(x) + C_1 \sin(q_1 x) + C_2 \sin(q_2 x) + \dots, \quad (19)$$

where the common multiples of $1, q_1, q_2, \dots$, are irrational numbers in pairs, the projective matrix becomes a vector

$$\mathcal{S} = (1, q_1, q_2, \dots). \quad (20)$$

Then our proposed method can be applied to calculate one-dimensional quasicrystals.

In the following, a lemma is given to indicate which variable should be computed in PM.

Lemma 1. *For a d -dimensional quasiperiodic function $g(\mathbf{r})$, under the expansion (17), we have*

$$\lim_{V \rightarrow \infty} \frac{1}{V} \int d\mathbf{r} g(\mathbf{r}) = \hat{g}(\mathbf{H}) \Big|_{\mathbf{H}=\mathbf{0}}. \quad (21)$$

Proof. Firstly, we note that

$$\lim_{V \rightarrow \infty} \frac{1}{V} \int d\mathbf{r} \exp \left\{ (\mathcal{S} \cdot \mathbf{H}) \cdot \mathbf{r} \right\} = \delta(\mathcal{S} \cdot \mathbf{H}). \quad (22)$$

Therefore,

$$\lim_{V \rightarrow \infty} \frac{1}{V} \int d\mathbf{r} g(\mathbf{r}) = \lim_{V \rightarrow \infty} \frac{1}{V} \int d\mathbf{r} \sum_{\mathbf{H}} \hat{g}(\mathbf{H}) e^{i[(\mathcal{S} \cdot \mathbf{H})^T \cdot \mathbf{r}]} = \hat{g}(\mathbf{H}) \Big|_{\mathcal{S} \cdot \mathbf{H} = 0}. \quad (23)$$

Then we just need to prove that the expressions (21) and (23) are equivalent. From the definition of (16), n -dimensional reciprocal vector $\mathbf{H}$ is the integer-valued combinations of linearly independent primitive reciprocal vectors $\mathbf{b}_i^*$, $i = 1, 2, \dots, n$. Because the projective matrix $\mathcal{S}$ is linear, the d -dimensional projected vector $\mathcal{S} \cdot \mathbf{H}$ is also the integer-valued combinations of the d -dimensional projected vectors $\mathbf{p}_i^* = \mathcal{S} \cdot \mathbf{b}_i^*$. $\mathcal{S} \cdot \mathbf{H} = 0$ is equivalent to the integral coefficients h_i in Eqn. (23) to be zero. It means that the n -dimensional reciprocal vector $\mathbf{H} = 0$. $\square$

Remark 1. From Lemma 1, the Fourier coefficients $\hat{g}(\mathbf{H})$ rather than $\hat{g}(\mathcal{S} \cdot \mathbf{H})$ should be computed in PM.

In PM, a quasicrystal is computed in n -dimensional reciprocal space as a periodic structure, then the n -dimensional structure is projected into d -dimensional space to obtain the d -dimensional quasicrystal. Since different periodic phases have their own periodicity, the appropriate computational box is important to determine the final morphology of solutions, especially for complex phases. For example, in diblock copolymer systems²⁵, the lamellae phase can be obtained easily in any computational region, however, for complex gyroid pattern, the computational box should be close to its period. For more complex quasicrystal, the computational box should be estimated carefully before computing. We also note that a equilibrium periodic structure is not only the minimum of a free energy density F with respect to order parameters, but also with respect to the unit cell²⁴. Therefore, the unit cell in n -dimensional space, $\mathcal{B} = [\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n]$, should satisfy $\partial F / \partial \mathcal{B} = 0$. We will give a estimation formula for the unit cell of quasicrystals in the Lifshitz-Petrich model (see Sec. IIB 3).

The PM is just with the help of n -dimensional reciprocal space to compute the d -dimensional quasicrystal. The physical space variable $\mathbf{r}$ always belongs to d -dimensional space. Therefore an energy functional including d -dimensional quasicrystals should not be

up to describing n -dimensional structures. The PM is able to compute quasicrystals rather than crystalline approximants. The PM is also using the periodic condition conveniently in higher-dimensional reciprocal space. Compared with CAM, the most important advantage of PM is that the method overcomes the restriction of SDA. Therefore PM can compute the free energy density to high accuracy numerically. Meanwhile, the proposed method can implement in a n -dimensional reciprocal unit cell to reduce computational cost.

2. An Example of 12-fold Rotational Symmetry

As an example, we take the 12-fold rotational symmetry in two dimensions to illustrate the PM. As Fig.2 shows, the 12 reciprocal lattice vectors cannot be represented by two




FIG. 3: Sets of reciprocal lattice vectors $\{\mathbf{H}\}$ of the 12-fold rotational symmetry with 4 components.

noncollinear vectors $\mathbf{e}_1^*$ and $\mathbf{e}_2^*$ with integral coefficients. However, as Fig.3 shows, if the vectors of $\mathbf{b}_1^*$, $\mathbf{b}_2^*$, $\mathbf{b}_3^*$, and $\mathbf{b}_4^*$ with 4 components are chosen as primitive reciprocal vectors, other reciprocal lattice vectors can be represented as integral combinations of these primitive vectors. The selected 4 vectors make up a basis in 4-dimensional space. The 2-dimensional 12-fold example is related to a periodic structure in 4-dimensional reciprocal space. The projected vectors $\mathbf{p}_i^* = \mathcal{S} \cdot \mathbf{b}_i^*$, $i = 1, 2, 3, 4$ also can be expressed by $\mathbf{e}_1^*$, $\mathbf{e}_2^*$ in 2-dimensional space, i.e.,

$$p_{i1}^* = \cos \frac{(i-1)\pi}{6} e_{11}^*, \quad p_{i2}^* = \sin \frac{(i-1)\pi}{6} e_{22}^*. \quad (24)$$

Therefore, the projective matrix is

$$\mathcal{S} = \begin{pmatrix} 1 & \cos(\pi/6) & \cos(\pi/3) & 0 \\ 0 & \sin(\pi/6) & \sin(\pi/3) & 1 \end{pmatrix}. \quad (25)$$

3. Applying PM to Lifshitz-Petrich Method

Applying PM to the Lifshitz-Petrich model, the order parameter $\phi(x, y)$ is expanded as

$$\phi(x, y) = \sum_{\mathbf{H}} \hat{\phi}(\mathbf{H}) \exp \left\{ i(\mathcal{S} \cdot \mathbf{H})^T \cdot (x, y)^T \right\}, \quad (26)$$

where $\mathbf{H}$ is a n -dimensional vector, $\mathcal{S}$ is a $2 \times n$ -order projective matrix. For quasicrystals, the system volume V should go to infinity. Based on Lemma 1, the Lifshitz-Petrich free energy density functional becomes

$$\begin{aligned} F[\phi(x, y)] = & \frac{1}{2} \sum_{\mathbf{H}_1 + \mathbf{H}_2 = 0} \{c[1 - (g_1^2 + g_2^2)]^2 [q^2 - (g_1^2 + g_2^2)]^2 - \varepsilon\} \hat{\phi}(\mathbf{H}_1) \hat{\phi}(\mathbf{H}_2) \\ & - \frac{\alpha}{3} \sum_{\mathbf{H}_1 + \mathbf{H}_2 + \mathbf{H}_3 = 0} \hat{\phi}(\mathbf{H}_1) \hat{\phi}(\mathbf{H}_2) \hat{\phi}(\mathbf{H}_3) + \frac{1}{4} \sum_{\mathbf{H}_1 + \mathbf{H}_2 + \mathbf{H}_3 + \mathbf{H}_4 = 0} \hat{\phi}(\mathbf{H}_1) \hat{\phi}(\mathbf{H}_2) \hat{\phi}(\mathbf{H}_3) \hat{\phi}(\mathbf{H}_4), \end{aligned} \quad (27)$$

where g_1 and g_2 are defined by

$$\mathcal{S} \cdot \mathbf{H} = \left(\sum_{i=1}^n s_{1i} \sum_{j=1}^n h_j b_{ji}, \sum_{i=1}^n s_{2i} \sum_{j=1}^n h_j b_{ji} \right)^T \triangleq (g_1, g_2)^T, \quad (28)$$

$h_j \in \mathcal{Z}$, s_{1j} and s_{2j} are the components of the projective matrix $\mathcal{S}$, and b_{ji} is the i -th component of the primitive reciprocal vector $\mathbf{b}_j$. The n -dimensional Fourier coefficient $\hat{\phi}(\mathbf{H})$ can be solved by the semi-implicit method (4)

$$\begin{aligned} & \left(\frac{1}{\Delta t} + c(1 - g_1^2 - g_2^2)^2 (q^2 - g_1^2 - g_2^2)^2 \right) \hat{\phi}_{t+\Delta t}(\mathbf{H}) \\ & = \left(\frac{1}{\Delta t} + \varepsilon \right) \hat{\phi}_t(\mathbf{H}) + \alpha(\phi_t^2)(\mathbf{H}) - (\phi_t^3)(\mathbf{H}), \end{aligned} \quad (29)$$

where the quadratic term and the third term are

$$(\phi_t^2)(\mathbf{H}) = \sum_{\mathbf{H}_1 + \mathbf{H}_2 = \mathbf{H}} \hat{\phi}_t(\mathbf{H}_1) \hat{\phi}_t(\mathbf{H}_2), \quad (\phi_t^3)(\mathbf{H}) = \sum_{\mathbf{H}_1 + \mathbf{H}_2 + \mathbf{H}_3 = \mathbf{H}} \hat{\phi}_t(\mathbf{H}_1) \hat{\phi}_t(\mathbf{H}_2) \hat{\phi}_t(\mathbf{H}_3). \quad (30)$$

In Eqn. (29), the linear term can be solved easily. The nonlinear terms are n -dimensional convolutions in reciprocal space. Directly computing will result in expansive computational

cost. However these convolutions are local point multiplication in n -dimensional direct space. Therefore we use the pseudospectral method²⁴ to treat these terms by FFT. It should be emphasized that the PM is implemented in n -dimensional reciprocal space than d -dimensional physical space. Computing these convolutions in n -dimensional direct space is to reduce computational complexity.

For PM, we give a method to estimate the n -dimensional unit cell for the Lifshitz-Petrich model²⁵. Without loss of generality, we can choose a proper coordinate system such that $b_{ii} \neq 0$, $b_{ij} = 0$, when $j > i$. If $\mathcal{B} = [\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n]$ is a n -dimensional unit cell of a d -dimensional quasicrystal. The first deviations of the free energy functional with respect to b_{ij} should be zero, where $i = 1, 2, \dots, n$, $j = 1, 2, \dots, i$, i.e.,

$$\sum_{\mathbf{H}} h_j (g_1 s_{1i} + g_2 s_{2i}) [1 - (g_1^2 + g_2^2)] [q^2 - (g_1^2 + g_2^2)] [1 + q^2 - 2(g_1^2 + g_2^2)] \cdot |\hat{\phi}(\mathbf{H})|^2 = 0. \quad (31)$$

Then the unit cell can be obtained by solving the Eqn. (31). Since the Fourier coefficients $\phi_{\mathbf{H}}$ are unknown beforehand, in practice, only primary reciprocal vectors with equal Fourier coefficients are considered in estimating the unit cell.

C. Computational Complexity

In this section, we will give a general analysis on the computational complexity of CAM and PM in solving the two-dimensional Lifshitz-Petrich model. The computational complexity of CAM is dependent on the approximate accuracy of SDA (11) and the numerical resolution. However, the PM overcomes the restriction of SDA, whose computational complexity is only dependent on the numerical resolution. As described in Sec II A 1 and Sec. II B 1, both CAM and PM can use pseudospectral method in computing the two-dimensional Lifshitz-Petrich model. Then the number of basic functions used in the reciprocal space is equivalent to the number of discretized points in the direct space. In order to guarantee the same numerical resolution, we assume that the mesh step size is Δx both for CAM (2 dimensions) and PM (n dimensions) in the direct space.

In the Lifshitz-Petrich model, the CAM is implemented in two dimensions. At a desired approximate error E_{SDA} as well as the integer L , the grid size is $N_{\mathbf{k}} \times N_{\mathbf{k}}$,

$$N_{\mathbf{k}} = \left\lceil \frac{D_{\mathbf{k}}}{\Delta x} \right\rceil, \quad D_{\mathbf{k}} = 2\pi \times L, \quad (32)$$

where $[\cdot]$ rounds the number $\cdot$ to the nearest integer. Then the computational complexity of CAM

$$O(N_{t,\mathbf{k}} \cdot 2N_{\mathbf{k}}^2 \log N_{\mathbf{k}}), \quad (33)$$

where $N_{t,\mathbf{k}}$ is the number of time iterations.

In PM, the 2-dimensional quasicrystal is represented in n -dimensional reciprocal space as a periodic structure. The dimension of n is dependent on the rotational symmetry of a quasicrystal. Therefore the 2-dimensional quasicrystal can be computed in a n -dimensional unit cell. We assume that the size of the n -dimensional unit cell is $D_{\mathbf{H}}$ in the direct space. Then the number of the basic functions is $N_{\mathbf{H}}^n$, with

$$N_{\mathbf{H}} = \left\lceil \frac{D_{\mathbf{H}}}{\Delta x} \right\rceil. \quad (34)$$

As discussed in Sec. II B 1, the pseudospectral method in PM is used to solve the dynamical equation (29) in n -dimensional reciprocal space with the help of FFT. Therefore the computational complexity of PM is

$$O(N_{t,\mathbf{H}} \cdot nN_{\mathbf{H}}^n \log N_{\mathbf{H}}), \quad (35)$$

where $N_{t,\mathbf{H}}$ is the number of time iterations.

From the estimation formulas (33), the computational complexity of CAM is a function of the integer L , which is dependent upon the approximate error of SDA. In next section, we will find that $D_{\mathbf{k}} \gg D_{\mathbf{H}}$ as the approximate error of SDA decreases. The computational complexity of CAM may be larger than that of PM in situation where higher numerical accuracy is required. More importantly, PM can compute quasicrystals and their free energy density to high accuracy without any approximate error of SDA. However, CAM always has the approximate error of SDA unless the computational box goes to infinity which results in unaccepted computational cost.

III. NUMERICAL RESULTS AND DISCUSSION

We will demonstrate the behavior of the two numerical methods, the CAM and the PM, based on the two-dimensional Lifshitz-Petrich model. In previous research^{22,26}, the Lifshitz-Petrich model showed that if q is chosen around $2 \cos(\pi/12)$ one can obtain a 2-dimensional

quasicrystal with dodecagonal (12-fold) symmetry, the earlier work found that no choice of q yields globally stable octagonal or decagonal symmetric pattern. However, in the recent study, the decagonal quasicrystal is found to be stable in the Lifshitz-Petrich model²⁷. Here we only consider the dodecagonal symmetric structure.

A. Computational Complexity of Dodecagonal Symmetric Structure

For the dodecagonal quasicrystal (DDQC), the initial reciprocal vectors at which the Fourier coefficients are nonzero. for DDQC in the Lifshitz-Petrich model is shown in Fig. 4, which contain two 12-fold stars of wave vectors, one on $|\mathbf{k}| = 1$ and other on $|\mathbf{k}| = q = 2 \cos(\pi/12)$ ²². The specific reciprocal vectors are given in Tab.I when $\mathbf{e}_1^* = (1, 0)$ and




FIG. 4: Initial reciprocal lattice vectors for DDQC.

TABLE I: Initial reciprocal lattice vectors for DDQC in 2-dimensional space with $q = 2 \cos(\pi/12)$.

$ \mathbf{k} = 1$	$(\cos(j\pi/6), \sin(j\pi/6)), j = 0, 1, \dots, 11$
$ \mathbf{k} = q$	$(q \cos(j\pi/6 + \pi/12), q \sin(j\pi/6 + \pi/12)), j = 0, 1, \dots, 11$

$\mathbf{e}_2^* = (0, 1)$, as shown in Fig2, are chosen as the basic reciprocal vectors in 2 dimensions. The distinct nonzero represented coefficients of these initial reciprocal vectors for DDQC phase are 1, $1/2$, $\sqrt{3}/2$, $2 \cos(\pi/12)$, $\cos(\pi/12)$ and $\sqrt{3} \cos(\pi/12)$. The condition of SDA (11) must be satisfied when one uses CAM. Fig. 6 shows the trend of the approximate error of




FIG. 5: The trend of the approximate error of SDA, E_{SDA} , as a function of the integer L .

SDA, E_{SDA} , as the integer L increases. From the Fig. 6, one can find that the convergence of E_{SDA} is not uniform. Tab.II gives the minimal integer for desired approximate error

TABLE II: The minimal integer L for desired approximate error of SDA, E_{SDA} .

E_{SDA}	0.19098	0.17486	0.07042	0.04953	0.03583	0.02961	0.01936
L	30	208	410	3404	6016	32312	82262

The size of computational box used in CAM is $D_{\mathbf{k}} = 2\pi L$.

E_{SDA} , and the corresponding size $D_{\mathbf{k}}$ of computational box used in CAM. Then the initial reciprocal vectors in CAM are $[L \cdot \mathbf{k}]$. From the table, we find that L increases quickly as the approximate error becomes small. Accordingly, the computational cost will increase greatly. We also find the $E_{SDA} \approx 0.19098$ ($L = 30$) is the least requirement for computing dodecagonal symmetric structure, it is consistent with the result in Ref. [22]. The initial reciprocal vectors in 4-dimensional space representation of PM are shown in Tab. III when the primitive reciprocal vectors are $\mathbf{b}_1^* = (1, 0, 0, 0)$, $\mathbf{b}_2^* = (0, 1, 0, 0)$, $\mathbf{b}_3^* = (0, 0, 1, 0)$, and $\mathbf{b}_4^* = (0, 0, 0, 1)$, as shown in Fig 3. We can use the 24 initial reciprocal vectors in Tab. III to estimate the unit cell of DDQC in 4-dimensional space. The projective matrix $\mathcal{S}$ of PM is given by Eqn. (25). From the estimation formula (31) and the dual relationship (6), the size of the unit cell in n -dimensional direct space is $D_{\mathbf{H}} = 2\pi$.

In the following, we will compare the computational complexity between CAM of different unit cell with $D_{\mathbf{k}}$ and PM as discussed in Sec. II C. In each time step, for the dodecagonal

TABLE III: Initial reciprocal lattice vectors $\{\mathbf{H}\}$ for DDQC in PM with $q = 2 \cos(\pi/12)$.

$ \mathcal{S} \cdot \mathbf{H} = 1$	(0 1 0 -1) (0 -1 0 1) (1 0 0 0) (-1 0 0 0) (0 1 0 0) (0 -1 0 0) (0 0 1 0) (0 0 -1 0) (0 0 0 1) (0 0 0 -1) (-1 0 1 0) (1 0 -1 0)
$ \mathcal{S} \cdot \mathbf{H} = q$	(1 1 0 -1) (-1 -1 0 1) (1 1 0 0) (-1 -1 0 0) (0 1 1 0) (0 -1 -1 0) (0 0 1 1) (0 0 -1 -1) (-1 0 1 1) (1 0 -1 -1) (-1 -1 1 1) (1 1 -1 -1)

The projective matrix $\mathcal{S}$ is given by Eqn. (25).

symmetric structure, the computational complexity of CAM is

$$C_{\mathbf{k}} = O(2N_{\mathbf{k}}^2 \log N_{\mathbf{k}}) = O\left(2\left(\frac{2\pi L}{\Delta x}\right)^2 \log\left(\frac{2\pi L}{\Delta x}\right)\right), \quad (36)$$

and the computational complexity of PM is

$$C_{\mathbf{H}} = O(4N_{\mathbf{H}}^4 \log N_{\mathbf{H}}) = O\left(4\left(\frac{2\pi}{\Delta x}\right)^4 \log\left(\frac{2\pi}{\Delta x}\right)\right). \quad (37)$$

For $L = 30$ with $E_{SDA} \approx 0.19098$ in CAM, the computational complexity $C_{\mathbf{k}} < C_{\mathbf{H}}$ when Δx is smaller than 0.20943951. However, for $L = 208$, $C_{\mathbf{k}} < C_{\mathbf{H}}$ only if $\Delta x > 500.6549$, which is much larger than the size of the unit cell $D_{\mathbf{H}}$. It cannot be implemented numerically. The computational complexity $C_{\mathbf{H}}$ of PM is always smaller than that of $C_{\mathbf{k}}$. For higher approximate accuracy of SDA with larger L , $C_{\mathbf{H}} < C_{\mathbf{k}}$, and total computational complexity of PM may be less than CAM.

Then we compare the total computational complexity including time iterations of CAM (with $L = 30$) and PM through numerical experiments. All the methods considered here have been implemented in C language. Fourier transforms are computed using the FFTW²⁸ package. The codes were run in the same dedicated computer, a Intel(R) Xeon(R) CPU E5450 @3.00GH memory 16 G under linux. The time step size, Δt , is always selected as 0.1 for both methods. To measure error we use l^∞ norm,

$$f_{CAM} = \max_{\mathbf{k}} \left\{ \left(\frac{\delta F}{\delta \phi} \right) (\mathbf{k}) \right\} \quad (38)$$

for CAM, and

$$f_{PM} = \max_{\mathbf{H}} \left\{ \left(\frac{\delta F}{\delta \phi} \right) (\mathbf{H}) \right\} \quad (39)$$

for PM. In the following simulations, the error tolerance is set as 1.0×10^{-8} .




FIG. 6: The total CPU time, the free energy density and the number of time step at the error 10^{-8} of CAM ($L = 30$) and PM, with different mesh step size Δx at a set of parameters: $c = 50$, $\varepsilon = 0.015$, $\alpha = 1.0$.


Fig. 6 shows the total CPU time, the free energy density, the number of time step at the desired error 10^{-8} , of CAM ($L = 30$) and PM at a set of parameters, $c = 50$, $\varepsilon = 0.015$, $\alpha = 1.0$, with different discrete resolution. The number of time step of CAM on different grids is same, $N_{t,k} = 848$. PM also has the same time step to achieve the the error 10^{-8} on different grids, $N_{t,k} = 427$. It may be that the accuracy of spacial discretization is enough for the calculated structure at the set of parameter. For the same reason, the free energy density calculated by PM is nearly the same on these grids, while the CAM has the equal free energy density. However, the number of time step of CAM, $N_{t,k}$ is always large than that of PM, $N_{t,H}$. As discussed in Sec. IIIB, it may be that PM keeps the rotational symmetry, while CAM dose not. Then PM can reach the same accuracy quickly. The free energy density computed by PM is lower than the CAM. It may result from the approximate error of SDA in CAM. As Fig6 shown, the cost of CPU time of PM is lower than that of CAM until the mesh step size Δx is less than about 0.15. It is different from the above analysis about the computational complexity in each time step because of the different time step in PM and CAM.

From these results, we find that with relative large mesh step size Δx , PM can obtain enough numerical accuracy with less computational cost than CAM ($L = 30$). CAM always


has approximant error unless $D_{\mathbf{k}} \rightarrow \infty$. And as discussed in the section, higher accuracy of approximant error will result in heavy computational burden. Therefore, PM has less computational complexity than CAM to high accuracy in computing these quasicrystals represented in 4 dimensions, such as dodecagonal symmetric structures.

B. Dodecagonal Symmetric Structure

The initial reciprocal coefficients, as mentioned in Sec. III A, at which the Fourier coefficients are nonzero are used as initial values to find equilibrium dodecagonal symmetric structures. Unless otherwise specified, the simulations of CAM are performed on a 256×256 grids with $L = 30$, and the simulations of PM are performed with $N_{\mathbf{H}}^4 = 32^4$ plane waves in the following. The projective matrix $\mathcal{S}$ of PM is given by Eqn. (25). We also find that the morphologies and the free energy density will not change with denser grids. The morphology of the dodecagonal approximant computed by CAM in physical space is given in Fig. 7(a)(a). It is a periodic structure. The DDQC calculated by PM is shown in Fig. 7(b)(b), which can be viewed as a part of the dodecagonal approximant. Both morphologies demonstrate the



(a)Dodecagonal crystalline approximant



(b)DDQC

FIG. 7: The morphologies of (a). Dodecagonal approximant computed by CAM and (b). DDQC calculated by PM at $\varepsilon = 0.015$, $\alpha = 1$, $c = 50$, $q = 2 \cos(\pi/12)$.

12-fold orientational symmetry in the physical space, at least locally. However, further analysis about the relationship of the Fourier coefficients will come to a different conclusion. From the rotational symmetry, each 12 reciprocal vectors with dodecagonal rotational sym-

metry on the ring $|\mathbf{k}| = 1$ ($|\mathcal{S} \cdot \mathbf{H}| = 1$), or $|\mathbf{k}| = q$ ($|\mathcal{S} \cdot \mathbf{H}| = q$), should have equal Fourier coefficients. Tab.IV shows the principle Fourier coefficients calculated by CAM. Tab.V gives these Fourier coefficients on principle reciprocal vectors calculated by PM. From these results, we can find the 12 Fourier coefficients calculated by CAM on each characteristic ring are not equal. In contrast, as Tab. V shown, PM can obtain the 12 equal Fourier coefficients

TABLE IV: The Fourier coefficients of the principle reciprocal vectors for dodecagonal crystalline approximant computed by CAM with $L = 30$ at $\varepsilon = 0.015$, $\alpha = 1$, $c = 150$, $q = 2 \cos(\pi/12)$.

$ \mathbf{k} = L$	(30,0) (-30,0) (0,30) (0,-30)	6.106618210e-02
	(26,15) (-26,-15) (15,26) (-15,-26)	5.928204525e-02
	(-15,26) (15,-26) (-26,15) (26,-15)	-
$ \mathbf{k} = [L \cdot q]$	(41,41) (-41,-41) (-41,41) (41,-41)	5.458000478e-02
	(56,15) (-56,-15) (15,56) (-15,-56)	5.683018766e-02
	(-15,56) (15,-56) (-56,15) (56,-15)	-

on each ring. We also find the same phenomenon at different parameter coordinates. For

TABLE V: The Fourier coefficients of the principle reciprocal vectors for DDQC computed PM at $\varepsilon = 0.015$, $\alpha = 1$, $c = 150$, $q = 2 \cos(\pi/12)$.

$ \mathcal{S} \cdot \mathbf{k} = 1$	(0 1 0 -1) (0 -1 0 1) (1 0 0 0) (-1 0 0 0)	5.856822141e-02
	(0 1 0 0) (0 -1 0 0) (0 0 1 0) (0 0 -1 0)	
	(0 0 0 1) (0 0 0 -1) (-1 0 1 0) (1 0 -1 0)	
$ \mathcal{S} \cdot \mathbf{k} = q$	(1 1 0 -1) (-1 -1 0 1) (1 1 0 0) (-1 -1 0 0)	5.855442187e-02
	(0 1 1 0) (0 -1 -1 0) (0 0 1 1) (0 0 -1 -1)	
	(-1 0 1 1) (1 0 -1 -1) (-1 -1 1 1) (1 1 -1 -1)	

CAM, we also use higher approximate accuracy $E_{SDA} \approx 0.07042$, with the computational box $D_{\mathbf{k}} \times D_{\mathbf{k}}$, $D_{\mathbf{k}} = 820\pi$. The numerical experiments are implemented on a 4096×4096 grids. We find that CAM can capture the dodecagonal approximant, however, it cannot obtain 12 equal Fourier coefficients on each ring as well as the above numerical experiments. Therefore, we come to a conclusion that PM can keep the non-crystallographic symmetry

to high accuracy, while CAM cannot.

C. Free Energy Density

We compare the free energy computed by CAM, PM and single-wave approximation (SWA). The SWA uses these principal Fourier vectors to approximate the free energy functional²⁹. It translates the energy functional into a single-variable or multi-variable function. Then the minimum of free energy can be obtained by minimizing the reduced energy function with these variables. The principal Fourier vectors of DDQC have been shown in Fig 4. The reduced free energy function of DDQC computed by SWA in the Lifshitz-Petrich model^{22,27} is ($C \rightarrow \infty$)

$$F_{12} = 99(\phi_1^4 + \phi_q^4) + 144(\phi_1^3 + \phi_1^3\phi_q) + 360\phi_1^2\phi_q^2 - 24(\phi_1^2\phi_q + \phi_1^2\phi_q) - 8(\phi_1^3 + \phi_q^3) - 6\varepsilon(\phi_1^2 + \phi_q^2)/\alpha^2, \quad (40)$$

$\phi_1, \phi_q \in \mathcal{R}$ stand for Fourier coefficients on the $|\mathbf{k}| = 1$ and $|\mathbf{k}| = q$ rings, respectively. The approximated minimum of DDQC can be obtained by minimizing the above equation with respect to ϕ_1 and ϕ_q . The result of SWA can be viewed as a approximate solution of DDQC




FIG. 8: (a). Free energy density calculated by PM and CAM, as a function of the penalty factor c , relative to that of SWA ($c \rightarrow \infty$), with $q = 2 \cos(\pi/12)$, $\alpha = 1.0$, $\varepsilon = 0.015$. (b). Corresponding interfacial energy computed by PM and CAM.

when $c \rightarrow \infty$. Fig. 8(a) shows the free energy density calculated by PM and CAM, as a function of the penalty factor c , relative to that of SWA ($c \rightarrow \infty$) with $q = 2 \cos(\pi/12)$,

$\alpha = 1.0$, $\varepsilon = 0.015$. The free energy density F_{CAM} computed by CAM is less than F_{SWA} obtained by SWA until the penalty factor c increases to about 10. Then F_{CAM} is larger than F_{SWA} as c increases. The reason is that the CAM cannot control the principal reciprocal vectors located on $|\mathbf{k}| = 1$ and $|\mathbf{k}| = q$ when the penalty factor c increases, as shown in Fig. 8(b). In other words, CAM cannot calculate the interfacial energy (Laplacian terms) accurately when c is large. However the role of the differential terms in the Lifshitz-Petrich model is to keep the interactions at two characteristic scales. It is one of the reasons that the Lifshitz-Petrich model is able to stabilize quasicrystals²². Correspondingly, the F_{CAM} diverges from the true free energy density. We have also used $L = 410$ which improves the approximate accuracy of SDA in CAM to observe the free energy density. It costed much computational amount on a 4096×4096 grids, however, the same problem appeared. In contrast, the free energy density F_{PM} calculated by PM is always smaller than F_{SWA} for all c and converges to the F_{SWA} as $c \rightarrow \infty$. It is because the more basic functions are used in simulations by PM than that of SWA. And the approximated space of PM is more precise than that of SWA. Meanwhile, PM can maintain its principle reciprocal vectors located on scales 1 and q which is consistent with the model, as shown in Fig. 8(b). Therefore PM can compute the free energy density for all parameters of high numerical accuracy.

IV. CONCLUSIONS AND OUTLOOK

In the article, we summary the CAM, and point out the advantage of using the periodic condition and the restriction condition of the SDA in this method. Then we propose a new numerical method, PM, which is implemented in higher-dimensional reciprocal space. The developed method overcomes the restriction of SDA and enables us view a quasicrystal as a periodic structure, and uses the periodic condition conveniently. The projection method can reduce the computational effort efficiently by computed quasicrystals in a higher-dimensional unit cell and using the pseudospectral method. By applying two methods to the Lifshitz-Petrich model, we analyze the computational complexity. The computational complexity of CAM is dependent on the approximate error and the numerical resolution, while that of PM is only dependent upon the numerical resolution. Specially, in computing dodecagonal symmetric pattern, PM has less computational complexity than that of CAM to high accurate. We also find that our approach can keep the rotational symmetry accurately, and more

significantly, the algorithm can calculate the free energy density without any approximate error of SDA. However, the dimensions of our computational space are dependent on the rotational symmetry of quasicrystals. For quasiperiodic structures with 5-, 8-, 10- and 12-fold symmetry, the computational space dimension is four. PM has accepted computational complexity. For a quasicrystal with 7-, 9-, and 18-fold quasicrystals, the dimension of computational space is up to six. Therefore, our future work will focus on the improvement of the computational efficiency of our proposed method. Applying the proposed method to study real physical systems is also one of the major tasks in our further research. Finally, we should point out that PM can be also applied to study general d -dimensional aperiodic structures¹⁶ whose Fourier spectrum consists of δ -peaks, $H = \sum_{i=1}^n h_i \mathbf{b}_i^*$, $h_i \in \mathbb{Z}$, of rank n ($n > d$) with basis vectors $\mathbf{b}_i^*$, $i = 1, 2, \dots, n$.

ACKNOWLEDGMENTS

The authors would like to acknowledge Dr. Robert A. Wickham for his help with the English grammar and useful advice. The work is supported by the National Science Foundation of China 21274005 and 50930003, and the China Postdoctoral Science Foundation 2011M500179.

REFERENCES

- ¹D. Shechtman, I. Blech, D. Gratias, and J.W. Cahn. Metallic phase with long-range orientational order and no translational symmetry. *Physical Review Letters*, 53(20):1951–1953, 1984.
- ²D. Levine and P.J. Steinhardt. Quasicrystals: a new class of ordered structures. *Physical Review Letters*, 53(26):2477–2480, 1984.
- ³R. Penrose. The role of aesthetics in pure and applied mathematical research. *Bulletin of the Institute of Mathematics and Applications*, 10:266, 1974.
- ⁴M. Gardner. Extraordinary nonperiodic tiling that enriches the theory of tiles. *Scientific American*, 236:110–119, 1977.
- ⁵W. Steurer. Twenty years of structure research on quasicrystals. part I. pentagonal, octagonal, decagonal and dodecagonal quasicrystals. *Zeitschrift für Kristallogra-*

- phie/International journal for structural, physical, and chemical aspects of crystalline materials*, 219(7/2004):391–446, 2004, and references therein.
- ⁶A.P. Tsai. Icosahedral clusters, icosahedral order and stability of quasicrystals—view of metallurgy. *Science and Technology of Advanced Materials*, 9:013008, 2008, and references therein.
- ⁷X. Zeng, G. Ungar, Y. Liu, V. Percec, A.E. Dulcey, and J.K. Hobbs. Supramolecular dendritic liquid quasicrystals. *Nature*, 428(6979):157–160, 2004.
- ⁸X. Zeng, R. Kieffer, B. Glettner, C. Nürnberger, F. Liu, K. Pelz, M. Prehm, U. Baumeister, H. Hahn, H. Lang, G. A. Gehring, C. H. M. Weber, J. K. Hobbs, C. Tschierske, and G. Ungar. Complex multicolor tilings and critical phenomena in tetraphilic liquid crystals. *Science*, 331(6022):1302, 2011.
- ⁹V. Percec, M.R. Imam, M. Peterca, D.A. Wilson, and P.A. Heiney. Self-assembly of dendritic crowns into chiral supramolecular spheres. *Journal of the American Chemical Society*, 131(3):1294–1304, 2009.
- ¹⁰V. Percec, M.R. Imam, M. Peterca, D.A. Wilson, R. Graf, H.W. Spiess, V.S.K. Balagurusamy, and P.A. Heiney. Self-assembly of dendronized triphenylenes into helical pyramidal columns and chiral spheres. *Journal of the American Chemical Society*, 131(22):7662–7677, 2009.
- ¹¹K. Hayashida, T. Dotera, A. Takano, and Y. Matsushita. Polymeric quasicrystal: Mesoscopic quasicrystalline tiling in abc star polymers. *Physical Review Letters*, 98(19):195502, 2007.
- ¹²D.V. Talapin, E.V. Shevchenko, M.I. Bodnarchuk, X. Ye, J. Chen, and C.B. Murray. Quasicrystalline order in self-assembled binary nanoparticle superlattices. *Nature*, 461(7266):964–967, 2009.
- ¹³S. Fischer, A. Exner, K. Zielske, J. Perlich, S. Deloudi, W. Steurer, P. Lindner, and S. Förster. Colloidal quasicrystals with 12-fold and 18-fold diffraction symmetry. *Proceedings of the National Academy of Sciences*, 108(5):1810, 2011.
- ¹⁴L. Bindi, P.J. Steinhardt, N. Yao, and P.J. Lu. Natural quasicrystals. *Science*, 324(5932):1306, 2009.
- ¹⁵A.I. Goldman and R.F. Kelton. Quasicrystals and crystalline approximants. *Reviews of Modern Physics*, 65:213–230, 1993.

- ¹⁶T. Janssen, G. Chapuis, and M. De Boissieu. *Aperiodic Crystals: From Modulated Phases to Quasicrystals*, volume 20. Oxford University Press, USA, 2007.
- ¹⁷W. Steurer and S. Deloudi. *Crystallography of Quasicrystals: Concepts, Methods and Structures*, volume 126. Springer Verlag, 2009.
- ¹⁸J. Bamberg, G. Cairns, and D. Kilminster. The crystallographic restriction, permutations, and goldbach’s conjecture. *The American Mathematical Monthly*, 110(3):202–209, 2003.
- ¹⁹D.A. Rabson, N.D. Mermin, D.S. Rokhsar, and D.C. Wright. The space groups of axial crystals and quasicrystals. *Reviews of Modern Physics*, 63(3):699, 1991.
- ²⁰N.D. Mermin. The space groups of icosahedral quasicrystals and cubic, orthorhombic, monoclinic, and triclinic crystals. *Reviews of Modern Physics*, 64(1):3, 1992.
- ²¹J. Dräger and N.D. Mermin. Superspace groups without the embedding: the link between superspace and Fourier-space crystallography. *Physical Review Letters*, 76(9):1489–1492, 1996.
- ²²R. Lifshitz and D.M. Petrich. Theoretical model for faraday waves with multiple-frequency forcing. *Physical Review Letters*, 79(7):1261–1264, 1997.
- ²³H. Davenport and K. Mahler. Simultaneous diophantine approximation. *Duke Mathematical Journal*, 13:105–111, 1946.
- ²⁴P. Zhang and X. Zhang. An efficient numerical method of Landau-Brazovskii model. *Journal of Computational Physics*, 227(11):5859–5870, 2008.
- ²⁵K. Jiang, C. Wang, Y. Huang, and P. Zhang. Discovery of new metastable patterns in diblock copolymers. Accepted by *Communications in Computational Physics*.
- ²⁶K. Barkan, H. Diamant, and R. Lifshitz. Stability of quasicrystals composed of soft isotropic particles. *Physical Review B*, 83(17):172201, 2011.
- ²⁷K. Jiang, A.C. Shi, and P. Zhang. Stability of dodecagonal and decagonal quasicrystals in Lifshitz-Petrich model. In preparation.
- ²⁸M. Frigo, and S. Johnson, FFTW: An adaptive software architecture for the FFT. *ICASSP Conference Proceedings*, 3:1381–1384, 1988.
- ²⁹P.M. Chaikin and T.C. Lubensky. *Principles of Condensed Matter Physics*. Cambridge University Press, 1995.